

## **Spatial Structure of the Decapeptide Val-Ile-Lys-Lys-Ser-Thr-Ala-Leu-Leu-Gly in Water and in a Complex with Sodium Dodecyl Sulfate Micelles**

Blokhin D., Efimov S., Klochkov A., Yulmetov A., Filippov A., Antzutkin O., Aganov A., Klochkov V.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

### **Abstract**

We have studied the spatial structure of the decapeptide Val-Ile-Lys-Lys-Ser-Thr-Ala-Leu-Leu-Gly in aqueous solution and in a complex with sodium dodecyl sulfate (SDS) micelles by  $^1\text{H}$  nuclear magnetic resonance (NMR) spectroscopy and two-dimensional (2-D) NMR spectroscopy (total correlation spectroscopy and nuclear Overhauser effect spectroscopy (NOESY)). The approach used to determine the decapeptide spatial structure was based on analysis of the  $^1\text{H}$ - $^{13}\text{C}$  residual dipolar couplings in the molecules partially aligned in lyotropic liquid crystalline media. Analysis of the interproton distances obtained from the 2-D NOESY NMR spectrum was used to reveal the spatial structure of the decapeptide in a complex with SDS micelles. Complex formation was confirmed by analysis of  $^1\text{H}$  chemical shifts in the NMR spectrum of the decapeptide and analysis of the signs and values of NOEs in a solution with SDS micelles. © 2011 Springer-Verlag.

<http://dx.doi.org/10.1007/s00723-011-0257-x>

---